

**A MODIFIED SCHWINGER'S FORMULA
FOR THE CASIMIR EFFECT**

M.V. Cougo-Pinto, C.Farina

*Instituto de Física-Universidade Federal do Rio de Janeiro
Rio de Janeiro, R.J., CEP - 21.845-970 - Brasil*

Antonio J. Seguí-Santonja

*Departamento de Física Teórica
Universidad de Zaragoza - Zaragoza 50009 - Spain*
(A short version will appear in the Letters in Math. Phys.)

Abstract

After briefly reviewing how the (proper-time) Schwinger's formula works for computing the Casimir energy in the case of "scalar electrodynamics" where the boundary conditions are dictated by two perfectly conducting parallel plates with separation a in the Z -axis, we propose a slightly modification in the previous approach based on an analytical continuation method. As we will see, for the case at hand our formula does not need the use of Poisson summation to get a (renormalized) finite result.

In a recent letter Julian Schwinger⁽¹⁾ has taught us how to obtain the Casimir energy from the Schwinger formula for the effective action in the proper-time representation⁽²⁾:

$$W^{(1)}(s_0) = -\frac{i}{2} \int_{s_0}^{\infty} \frac{ds}{s} \text{Tr} e^{-isH} + \text{const.}, \quad (1)$$

where the exponential of $iW^{(1)}$ is defined as the vacuum persistency probability amplitude $\langle 0_+ | 0_- \rangle$ at the one-loop level, Tr means the total trace, H is the corresponding proper-time Hamiltonian and s_0 is a regularization cutoff that must be sent to zero only at the end of the calculations, after an appropriate subtraction of divergent terms is made; the additional constant is used to subtract divergent terms, thereby establishing the physical normalization for the situation.

Let us briefly show how equation (1) can be used to yield the correct Casimir energy in a simple example. Consider the situation of two perfectly conducting parallel plates with separation a along the Z -axis. In this case we have

$$H = [P_x^2 + P_y^2 - \omega^2] + P_z^2, \quad (2)$$

where $\vec{P} = -i\nabla$ and $\omega = +i\partial_t$. Observing that, while the operators in the bracket on the r.h.s. of (2) have continuous spectra, the operator P_z^2 has a discretized one, given by $p_z^2 = \left(\frac{n\pi}{a}\right)^2$; $n = 1, 2, \dots$, a straightforward calculation shows that

$$\text{Tr} e^{-isH} = AT \frac{1}{\sqrt{i(4\pi s)^3}} \sum_{n=1}^{\infty} e^{-is\left(\frac{n\pi}{a}\right)^2}, \quad (3)$$

where A is the (finite, although as big as we want) area of each plate and T the finite duration of the measurement.

After identifying the vacuum energy shift $\mathcal{E}$ with $-\frac{W^{(1)}}{T}$, we obtain from (1) and (3) that

$$\frac{\mathcal{E}(s_0; a)}{A} = \frac{\sqrt{i}}{16\pi^{3/2}} \int_{s_0}^{\infty} \frac{ds}{s^3} \sum_{n=1}^{\infty} e^{-in^2 s (\pi/a)^2}. \quad (4)$$

It is obvious that the limit $s_0 \rightarrow 0$ gives a divergent result so that we need to separate the infinities from the relevant finite part of equation (4). With this goal, we shall use the Poisson summation formula,

$$\sum_{n=-\infty}^{\infty} e^{-n^2 \pi \tau} = \sqrt{\frac{1}{\tau}} \sum_{n=-\infty}^{\infty} e^{-n^2 \pi (1/\tau)}. \quad (5)$$

It follows from (5) that

$$\sum_{n=1}^{\infty} e^{-in^2 s(\pi/a)^2} = \frac{a}{2} \frac{1}{\sqrt{i\pi s}} - \frac{1}{2} + \frac{a}{\sqrt{i\pi s}} \sum_{n=1}^{\infty} e^{i(na)^2/s}. \quad (6)$$

Inserting (6) into (4), we have

$$\begin{aligned} \frac{\mathcal{E}(s_0; a)}{A} &= \frac{1}{A} (Aa) \frac{1}{32\pi^2} \int_{s_0}^{\infty} \frac{ds}{s^3} - \frac{1}{A} A \frac{\sqrt{i}}{32\pi^{3/2}} \int_{s_0}^{\infty} \frac{ds}{s^{5/2}} + \\ &+ \frac{a}{16\pi^2} \int_{s_0}^{\infty} \frac{ds}{s^3} \sum_{n=1}^{\infty} e^{i(na)^2/s}. \end{aligned} \quad (7)$$

The first two terms on the r.h.s. of (7) are divergent when the limit $s_0 \rightarrow 0$ is taken, and hence they must be discarded. The physical reason for that is the following: the first one is proportional to the spatial volume Aa between the two plates, giving a uniform (infinite) contribution for the Casimir energy which is independent of the positions of the plates. Since this term violates the correct normalization of $\mathcal{E}$ ($\mathcal{E} = 0$ for $a \rightarrow \infty$), we must absorb it in the normalization of $\langle 0_+ | 0_- \rangle$. The second one does not contain a , so that it corresponds to the (infinite) energy required to create both plates; since we are only interested in energy shifts caused by changing the relative configuration of the plates we must also disregard this contribution.

Therefore, we are left with

$$\begin{aligned} \frac{\mathcal{E}(a)}{A} &= \lim_{s_0 \rightarrow 0} \frac{a}{16\pi^2} \int_{s_0}^{\infty} \frac{ds}{s^3} \sum_{n=1}^{\infty} e^{i(na)^2/s} \\ &= -\frac{1}{16\pi^2} \frac{1}{a^3} \sum_{n=1}^{\infty} \int_0^{\infty} d\sigma \sigma e^{-n^2 \sigma}, \end{aligned} \quad (8)$$

where we made the convenient change of variables $\sigma = \frac{a^2}{is}$.

Appealing to the well known formulas

$$\int_0^{\infty} dz z^{\nu-1} e^{-Az} = \Gamma(\nu) A^{-\nu} \quad (Re(\nu) > 0), \quad (9a)$$

$$\zeta_R(s) = \sum_{n=1}^{\infty} n^{-s} \quad (Re(s) > 1), \quad (9b)$$

where $\zeta_R(s)$ is the Riemann zeta function, we get

$$\begin{aligned}\frac{\mathcal{E}(a)}{A} &= -\frac{1}{16\pi^2} \frac{1}{a^3} \zeta_R(4) \\ &= -\frac{\pi^2}{1440} \frac{1}{a^3},\end{aligned}\tag{10}$$

which, apart from the factor 2 that takes into account the two photon polarizations (this simple multiplication by 2 does not work when other geometries are involved), is the correct result found by other methods⁽³⁾.

Now, we shall show that the Schwinger's method for a massless scalar theory, with a slight modification in the regularization process, leads to the finite result in an exceedingly short and simple way, without requiring any subtraction of infinities. The appearance of any divergent term during the calculation is bypassed by means of analytical continuation of the functions involved.

The basic idea of our approach is the following: since the origin of the divergent result in (1) when taking $s_0 \rightarrow 0$ was the presence of negative powers of s in the integrand, we can modify equation (1) by including as many positive powers of s as necessary (in fact we shall include the factor s^ν) so that we will be able to release the lower limit of the integral to zero. Therefore, our starting point is given by

$$W^{(1)}(\nu) = -\frac{1}{2} \int_0^\infty \frac{ds}{s} s^\nu \text{Tr} e^{-isH},\tag{11}$$

where ν must be sufficiently large to regularize the integral. After the integral is evaluated, we make an analytical extension of $W^{(1)}(\nu)$ to the whole complex plane (of ν) and then take the limit $\nu \rightarrow 0$. As we will see, for the case at hand, this procedure will give directly the desired (finite) result.

Inserting (3) into (11), we have (after identifying $\mathcal{E} = -\frac{W^{(1)}}{T}$)

$$\begin{aligned}\frac{\mathcal{E}(\nu; a)}{A} &= \frac{\sqrt{i}}{16\pi^{3/2}} \sum_{n=1}^\infty \int_0^\infty ds s^{\nu-3/2-1} e^{-s(i\pi^2 n^2/a^2)} \\ &= -\frac{\pi^2}{16} \frac{1}{\sqrt{\pi}} \Gamma(\nu - 3/2) \zeta_R(2\nu - 3) \frac{1}{a^3},\end{aligned}\tag{12}$$

where we used equations (9a) and (9b). To finish the calculation we take the limit $\nu \rightarrow 0$ and substitute the value of $\Gamma(-3/2)\zeta_R(-3)$. The result is

$$\frac{\mathcal{E}(a)}{A} = -\frac{\pi^2}{1440} \frac{1}{a^3},\tag{13}$$

in complete agreement with (10).

Let us finish this paper with two final remarks. First, that this last method works as well (with no need of subtracting infinities) in arbitrary $d + 1$ -dimensional space-time, leading us quickly to the well known result⁽³⁾:

$$\frac{\mathcal{E}}{A} = -\frac{1}{2}(\sqrt{\pi}/2)^d \Gamma(-\frac{d}{2}) \zeta_R(-d) \frac{1}{a^d}. \quad (13)$$

However, attention must be paid to the cases of even spatial dimensions (even d), where poles of the gamma function are cancelled by the zeros of the Riemann zeta function. Second, if our regularization, equation (11), method is applied to other theories, as QED with constant external fields, or even in the computation of the Casimir energy for a massive scalar field, subtraction of infinities may also be required before the analytical continuation to $\nu = 0$ is made. In a more detailed analysis, to be reported elsewhere, we will further explore these remarks.

We would like to thank to M. Asorey for a critical reading of the manuscript. This work was partially supported by MEC-CAICYT and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). One of us (A.J.S.S.) was also supported by DGI-CYT(Spain), grant PB90-0916.

References

1. Schwinger, J., *Lett. Math. Phys.* **24**, 59 (1992).
2. Schwinger, J., *Phys. Rev.* **82**, 664 (1951).
3. Plunien, G., Muller, B. and Greiner, W., *Phys. Rep.* **134**, 87 (1986).